


The NAoPri reports on all people (women and men) aged 18 and over newly diagnosed with primary breast cancer (stages 0 to 3) in NHS hospitals in England and Wales.



Triple Diagnostic Assessment

55% of people in England and 57% in Wales were reported to have Triple Diagnostic Assessment in a single hospital visit.





Surgery

86% of people in England and 86% in Wales received surgery within 12 months of diagnosis (stage 0 to stage 3A).





Breast Conserving Surgery (BCS)

72% of women in England and 68% in Wales had BCS. Mastectomy rates were higher with increased tumour size and older age.





Breast Reconstruction

24% of women in England and 14% in Wales had an immediate breast reconstruction following a mastectomy.





Chemotherapy

13% of people in England and 9% in Wales received neo-adjuvant chemotherapy (chemotherapy before surgery).

Among those with Early Invasive Breast Cancer (EIBC) having surgery, 33% of people in England and 29% in Wales received chemotherapy either before or after surgery.

Radiotherapy after BCS


85% of women in England with EIBC and 70% in Wales received adjuvant radiotherapy following BCS.





CNS Contact

For those with data available 98% of people in England and 99% in Wales had contact with a Clinical Nurse Specialist (CNS) after diagnosis. However, data completeness for England was 76%.



Survival

Percent of people who survived for 1 or 3 years after diagnosis in England and Wales (combined).

