


National Pancreatic Cancer Audit State of the Nation Report 2025: Methodology Supplement

An audit of care received by people diagnosed with pancreatic cancer between 1 January 2021 to 31 December 2022 in England and 1 January 2022 and 31 December 2023 in Wales.

Published September 2025; updated November 2025



Citation for this document:

National Pancreatic Cancer Audit (NPaCA) State of the Nation Report 2025. London:
National Cancer Audit Collaborating Centre, Royal College of Surgeons of England, 2025.

This document was prepared by members of the NPaCA project team:

Ganesh Radhakrishna, Co-Clinical Lead/Consultant Clinical Oncologist
Andrew Smith, Co-Clinical Lead/Consultant Surgeon
Nigel Trudgill, Co-Clinical Lead, Consultant Gastroenterologist
David Cromwell, Senior Methodologist/Professor of Health Services Research
Vikki Hart, Senior Project Manager
Amanda McDonnell, Data Scientist
Suzi Nallamilli, Clinical Fellow
Min Hae Park, Methodologist/Assistant Professor
Faine Chan, Project Coordinator

With review and input from:

[NPaCA Clinical Reference Group](#)

[NATCAN Executive Team](#)



The Royal College of Surgeons of England is an independent professional body committed to enabling surgeons to achieve and maintain the highest standards of surgical practice and patient care. As part of this it supports audit and the evaluation of clinical effectiveness for surgery. Registered Charity no: 212808.



The National Cancer Audit Collaborating Centre (NATCAN) is commissioned by the [Healthcare Quality Improvement Partnership \(HQIP\)](#) and funded by NHS England and the Welsh Government as part of the [National Clinical Audit and Patient Outcomes Programme](#) (NCAPOP). NATCAN delivers national audits in bowel, breast (primary and metastatic), kidney, lung, non-Hodgkin lymphoma, oesophago-gastric, ovarian, pancreatic and prostate cancers.



The Association of Upper Gastrointestinal Surgery of Great Britain and Ireland is the speciality society that represents upper gastrointestinal surgeons. It is one of the key partners leading the Audit. Registered Charity no: 1093090



British Society of Gastroenterology is the speciality society of gastroenterologists. It is one of the key partners leading the Audit. Registered Charity no: 1149074



Royal College of Radiologists is the professional body for clinical radiologists and clinical oncologists. It is one of the key partners leading the Audit. Registered Charity no: 211540



This work uses data that has been provided by patients and collected by the NHS as part of their care and support. For patients diagnosed in England, the data is collated, maintained and quality assured by the National Disease Registration Service (NDRS), which is part of NHS England. Access to the data was facilitated by the NHS England Data Access Request Service.



NHS Wales is implementing a new cancer informatics system. As a result, the quality and completeness of data from Wales is likely to have been impacted due to implementation of this new system across multiple NHS organisations (Health Boards), which has resulted in data being supplied by both old and new systems. Additionally, and reflecting the uncertainty of data quality, the data submitted to the audit may not have undergone routine clinical validation prior to submission to the Wales Cancer Network (WCN), Public Health Wales.

Contents

1. Introduction.....	3
2. Sources of Data.....	4
3. Inclusion and Exclusion Criteria.....	4
4. Key Data Items.....	6
5. Indicator Definitions.....	12
5.1 Performance Indicator 1: FDG-PET/CT scan prior to surgery.....	12
5.2 Performance Indicator 2: Multidisciplinary team (MDT) meeting.....	13
5.3 Performance Indicator 3: Biliary stent prior to surgery	14
5.4 Performance Indicator 4: Time to treatment.....	14
5.5 Performance Indicator 5: Receipt of disease-targeted treatment (stage 1-3).....	15
5.6 Performance Indicator 6: Receipt of disease-targeted treatment (stage 4)	16
5.7 Performance Indicator 7: Receipt of chemotherapy and/or radiotherapy alongside surgery.....	16
5.8 Performance Indicator 8: Clinical nurse specialist (CNS) involvement	17
5.9 Performance Indicator 9: Pancreatic enzyme replacement therapy (PERT) use	18
5.10 Performance Indicator 10: Survival from diagnosis	18
6. NHS organisations	19
7. Statistical Analysis	20
7.1 Small number Suppression	20
7.2 Risk-adjustment of indicators	20
7.3 Handling of missing data.....	21
8. Outlier Process	22
Appendix 1: Routine data sources.....	23
Appendix 2: Charlson Comorbidity Index.....	24
Appendix 3: Code lists	25
Appendix 4: AJCC TNM staging of (exocrine) pancreatic cancer.....	30
Appendix 5: World Health Organisation Performance Status	30

1. Introduction

This document provides supporting material to the 2025 State of the Nation (SotN) Report for the National Audit of Pancreatic Cancer (NPaCA) and its data tables and data viewer. The document describes the data used in the report with details on sources of data, criteria for inclusion and how data completeness, patient characteristics and performance indicators are derived and reported.

2. Sources of Data

The audit uses information from routine national health care datasets in England and Wales. These datasets capture details on the diagnosis, management, treatment and outcome of every patient newly diagnosed with cancer in the NHS in England and Wales.

For England, the audit cohort is based on the “Gold Standard” National Cancer Registration Data (NCRD) which is curated by the National Disease Registration Service (NDRS). The information held in the NCRD is compiled from a variety of sources including the Cancer Outcomes and Services Dataset (COSD), Hospital Episode Statistics admitted patient care (HES APC) records, the Systemic Anti-Cancer Therapy dataset (SACT), RTDS and data submitted by pathology laboratories. The audit also linked information from several other routine national health care datasets: see Appendix 1: Routine data sources for more detail on the data sources analysed as part of the audit.

As with cancer registries in other countries, cancer registrations in England can take up to 5 years after the end of a given calendar year to reach approximately 100% completeness and stability. NDRS uses an active system of gathering information on cancer diagnoses from multiple sources across the patient pathway. Completeness varies by tumour type because different patient pathways provide different opportunities for data flows into NDRS. The “Gold standard” cancer registration dataset that is used in cancer statistics bulletins and available for analysis outside of NDRS contains over 98% of all the people that will eventually be found by the registration process, and the completeness for a calendar year of data increases over time. More information about the cancer registration process can be found [here](#) and the timeliness of the NCRD [here](#).

The English data received by the National Cancer Audit Collaborating Centre (NATCAN) included data on people diagnosed with cancer up to 31st December 2022.

For Wales, the audit was provided with a registration dataset at patient level for patients diagnosed with cancer in 1st January 2022 to 31st December 2023. Welsh cancer registration data is captured through a national system, Cancer Information System for Wales (CaNISC) and the new Welsh Clinical Portal. The audit also received linked datasets of records from the Patient Episode Database for Wales (PEDW) containing information on inpatient and day case activity, and mortality data from the Office for National Statistics (ONS).

English and Welsh data were managed and analysed separately due to the different reporting periods.

3. Inclusion and Exclusion Criteria

The data submitted by NDRS and WCN is checked and filtered for eligible participants, tables 3.1 and 3.2 explain the process in defining the final cohort to be used in the audit.

People were included for analysis within the SotN Report if they met the following inclusion and not the exclusion criteria:

Table 3.1: Audit Inclusion Criteria	
Inclusion Criteria	Details
Type of cancer	Malignant neoplasm of the pancreas; Malignant neoplasm of the extrahepatic bile duct or ampullary tumours. See diagnosis codes listed in Appendix 3: Table 1
Adults	Age >=18

First Diagnosis	Earliest diagnosis date in the extract of data received <u>England</u> In instances of multiple pancreatic cancer diagnoses on the same day, prioritised records based on: • Worst stage, then • Record with most complete information across variables <u>Wales</u> • Only included patients who had a patient ID • Only included patients who had a 'date of diagnosis' (otherwise we are unable to confirm that they belong in this cohort) • Within duplicated patient rows, we prioritised those who had a diagnosis date, followed by those who had TNM staging detail, followed by most complete treatment information
Valid Diagnosis Date	England: 1st January 2021 to 31st December 2022 Wales: 1st January 2022 to 31st December 2023

Table 3.2: Audit Exclusion Criteria	
Exclusion Criteria	Details
Neuroendocrine tumours of the pancreas	For English data: In NCRD: Tumour site = C254 (Endocrine pancreas) and/or histology_coded_desc contains word "neuroendocrine" or "carcinoid" and/or morph_icd10_o2 or morph_coded contain neuroendocrine morphology codes specified in Appendix 3: Table 2 In SACT: morphology_clean contains neuroendocrine morphology codes specified in Appendix 3: Table 2 and/or benchmark_group = lanreotide or octreotide and/or drug_group = lanreotide or octreotide and primary_diagnosis="C24" or "C25" For Welsh data: Using CaNISC: if MORPHOLOGY_DESCRIPTION contains word "Carcinoid" or "Neuroendocrine" and/or TUMOUR_SITE_ICD10_CODE (CaNISC) = C25.4 and/or MORPHOLOGY_CODE contains neuroendocrine morphology codes specified in Appendix 3: Table 2
Reported by death certificate only or date of diagnosis corresponds to date of death	For English data: Using NCRD: final_route = DCO (Death Certificate Only) and/or basisofdiagnosis = 0 (Death certificate) and/or dco = Y (tumour registered from a death certificate only) and/or diagnosisdatebest = deathdatebest For Welsh data: DIAGNOSIS_DATE (CaNISC) or date_of_death (ONS) = date of death Note: Date of death derived from the earliest date of two variables: DATE_OF_DEATH (CaNISC) and date_of_death (ONS)

Diagnosed outside country of interest without treatment of interest	For English data: Trust of diagnosis was a Welsh health board (code starting with 7) and No record of major resection in England (Note: Trust code starting with “R” is in England) Note on Wales: If a patient was diagnosed in Wales, they have been included in the main cohort. If they were diagnosed in Wales, but part of their treatment pathway included an English hospital (eg. surgery in England), these they have still been included in the overall national figures, but they are removed from the denominator for surgical-related metrics.
---	--

4. Key Data Items

Details of the variables and datasets used to compile the data completeness are shown below in Table 4.1

Table 4.1: Data Completeness Variables				
Data Item	Source			
	England		Wales	
	Data field	Dataset	Data field	Dataset
Stage at Diagnosis	stage_best	NCRD	Derived using 3 variables T_STAGE_Final_Pretreatment, N_STAGE_Final_Pretreatment and M_STAGE_Final_Pretreatment to generate overall stage using the AJCC (American Joint Committee on Cancer staging) staging for pancreatic cancer version 8¹ – see Appendix 4: AJCC TNM staging of (exocrine) pancreatic cancer <i>Note on non-metastatic vs metastatic disease:</i> If the overall stage is unknown (eg. Missing T and N stage) but the M stage is 0, then the person is considered to be non-metastatic. Hence, more patients are grouped into the ‘non-metastatic’ cohort than reflected in the individual breakdown of staging groups.	CaNIS

¹ MB Amin, SB Edge, FL Greene, et al, eds. AJCC Cancer Staging Manual. 8th ed. New York: Springer; 2017.

Performance status	performancestatus	COSD	PERFORMANCE_STATUS	CaNISC
CNS (Clinical Nurse Specialist) involved	Derived using clinicalnursespecialist, counting any "Yes" response option as CNS involved	Derived by NDRS from COSD	Data not provided	

Details of the variables and datasets used to compile the patient and tumour characteristics are shown below in Table 4.2.

Table 4.2: Patient and Tumour Characteristics Variables				
<u>Data Item</u>	<u>Source</u>			
	England		Wales	
	<u>Data field</u>	<u>Dataset</u>	<u>Data field</u>	<u>Dataset</u>
Age at diagnosis	Age, categorised into 10 year groups	NCRD	Derived using the age at the start of the hospital episode closest to the date of diagnosis (EpisodeStartDate and PatientEpisodeStartAge Years, from PEDW)	PEDW
Diagnosis date	diagnosisdatebest Replaced diagnosis date with MDT meeting date when diagnosis date was equal to date of surgery. Note: not always possible as there were people with missing MDT meeting date	NCRD	DIAGNOSIS_DATE	CaNISC
Ethnicity	ethnicity Grouped as: White = 0, A, B, or C Mixed = D, E, F, or G Asian or Asian British = H, J, K, L, or R Black or Black British = M, N, or P Other ethnic group = S Missing = any other entry	NCRD	EthnicGroup	PEDW
Index of multiple deprivation	imd19_quintile_isoas	NCRD	QuintileGroup For duplicated patient rows in the LSOA dataset, with varying Quintiles, the lower quintile observation (i.e the most deprived) was used.	LSOA
Sex	gender	NCRD	Sex	PEDW

Performance status at diagnosis	performancestatus	COSD	PERFORMANCE_STATU S	CaNISC
Stage at Diagnosis	stage_best	NCRD	Derived using 3 variables T_STAGE_Final_Pretreatment, N_STAGE_Final_Pretreatment and M_STAGE_Final_Pretreatment to generate overall stage using the AJCC (American Joint Committee on Cancer staging) staging for pancreatic cancer version 8² – see Appendix 4: AJCC TNM staging of (exocrine) pancreatic cancer <i>Note on non-metastatic vs metastatic disease:</i> If the overall stage is unknown (eg. Missing T and N stage) but the M stage is 0, then the person is considered to be non-metastatic. Hence, more patients are grouped into the 'non-metastatic' cohort than reflected in the individual breakdown of staging groups.	CaNISC
Tumour site	Site_icd10	NCRD	TUMOUR_SITE_ICD10_CODE	CaNISC

Details of the variables and datasets used to construct performance indicators are listed below in Table 4.3.

Table 4.3 Indicator construction variables				
Data Item	Source			
	England		Wales	
	Data field	Dataset	Data field	Dataset
<i>Diagnosis, staging, and treatment planning</i>				
Biliary stent	Derived by searching variables opertn_1 – opertn_24 for biliary stent codes listed in Appendix 3: Table 5, up to 30 days before diagnosis date	HES-APC HES-OP	Derived by searching variables operation01 – operation12 for biliary stent codes listed in Table 5, up to 30 days before diagnosis date	PEDW
Biliary stent date	Earliest of opdate_01 – opdate_24 (HES-APC) or apptdate (HES-OP) associated with biliary	HES-APC HES-OP	Corresponding procedure dates taken from operation01datestyleddmmyy - operation12datestyle, up to 30 days before diagnosis date	PEDW

² MB Amin, SB Edge, FL Greene, et al, eds. AJCC Cancer Staging Manual. 8th ed. New York: Springer; 2017.

	stent record, up to 30 days before diagnosis date			
Imaging record	Derived by searching variable <code>imagingcodesnomedct</code> for imaging codes listed in Appendix 3: Table 3	DIDs	Patients who had a relevant scan were identified by the presence of a date in the variables: <code>Imaging__PET</code>	CaNISC
Imaging date	<code>Diagnostictestdate</code> associated with imaging record	DIDs	<code>Imaging__PET</code>	CaNISC
MDT meeting record / date	<code>firstmdtmeetingdate</code>	Derived by NDRS from COSD	Data not provided	
Organisation of diagnosis (Trust or local health board)	<code>diag_trust</code>	NCRD	<code>ORGANISATION_CODE</code>	CaNISC
Diagnosis date	<code>diagnosisdatebest</code> Replaced diagnosis date with MDT meeting date when diagnosis date was equal to date of surgery. Note: not always possible as there were people with missing MDT meeting date	NCRD	<code>DIAGNOSIS_DATE</code>	CaNISC
<i>Time from referral to start of treatment</i>				
Referral date	<code>crtp_date</code> Data cleaning: replaced to missing any referral dates more than 183 days (6 months) before diagnosis date or more than 7 days after diagnosis date In instances of multiple records per patient, executed the following steps to select referral date: - Dropped records if duplicates in terms of: patient ID, referral date, referral source, priority type, and site ICD10 code - Sorted records based on patient ID and referral date. In instances of duplicates in terms of referral date, prioritised records in the order of: (1) Containing an audit-eligible tumour site ICD-10 code (C25 or C24), (2) Complete data on referral source and	CWT	<code>DATE_OF_REFERRAL</code> Data cleaning: replaced to missing any referral dates more than 183 days (6 months) before diagnosis date or more than 7 days after diagnosis date	CaNISC

	priority type, (3) GP referral source , (4) Urgent priority referral - Selected record with earliest referral date, dropping other records for the patient			
Referral source	ref_source Grouped as follows: <i>GP referral</i> : 3 - "General medical practitioner" or 12 - "General practitioner with extended role"	CWT	SOURCE_OF_REFERRAL ("Referral from General Medical Practitioner" in the variable SOURCE_OF_REFERRAL)	CaNISC
Referral priority	priority_type Grouped as follows: <i>Urgent referral</i> : 2 – Urgent or 3 – Two Week Wait	CWT	Data not provided	
Treatment date	Derived as date of first record of disease-targeted treatment (see below)	Various – see below	Derived as date of first record of disease-targeted treatment (see below)	Various – see below
<i>Disease-targeted treatment</i>				
Surgery record	Derived by searching variable opcs4_code (NCRD) and opertn_1 – opertn_24 (HES-APC) for surgery codes listed in Appendix 3: Table 4	NCRD HES-APC	1. From PEDW: Derived by searching the variables operation01 – operation12 in PEDW for surgery codes listed in Table 4. 2. From CaNISC: Derived by searching for the surgery codes in the variable PROCEDURE_CODE	PEDW, CaNISC
Surgery date	eventdate (NCRD) or opdate_01 – opdate_24 (HES-APC) associated with surgery record	NCRD HES-APC	1. From PEDW: operation01datestyleddmmyyy – operation12datestyle 2. From CaNISC: DATE_OF_SURGERY Nb: In cases of multiple surgery dates identified, or discordance between datasets, the earliest date of surgery was used.	PEDW, CaNISC
Trust of surgery	trust_code (NCRD) or procode3 (HES-APC) associated with surgery record	NCRD HES-APC	1. From PEDW: ProviderOrganisationCode 2. From CaNISC data: PLACE_OF_SURGERY_CODE If the 2 datasets agreed on the date of the surgery, but there was a discrepancy in the location of the surgery location, then the location in the CaNISC dataset was used (as it has been more curated)	PEDW, CaNISC
SACT (Systemic Anti-Cancer Treatment)	Derived based on any record of anti-cancer treatment (except exclusions in Appendix 3: Table 6) in SACT and/or searching opertn_1 – opertn_24 (HES) for the chemotherapy administration codes in Appendix 3: Table 8	SACT HES-APC HES-OP	Derived based on a record of SACT in: • PEDW: Derived by searching the variables operation01 – operation12 in PEDW for SACT codes listed in Appendix 3 Table 8 and only SACT associated with a C24 or C25 diagnosis code in the variables Diagnosis01- Diagnosis14 were used. • CaNISC: the presence of a value in the variable START_DATE_OF_CHEMOTHERAPY • SACT dataset: the presence of a value in the variable DateofActivity.	PEDW, CaNISC , SACT dataset

			Nb: SACT dataset provided for people diagnosed in the 2023 calendar year only	
SACT date	Earliest of start_date_of_cycle (SACT) or opdate_01 – opdate_24 (HES-APC), or apptdate (HES-OP) associated with SACT treatment record	SACT	• PEDW: operation01datestyleddmmyyy – operation12datestyle • CaNISC: START_DATE_OF_CHEMOTHERAPY • SACT dataset: DateofActivity Nb: SACT dataset provided for people diagnosed in the 2023 calendar year only	PEDW, CaNISC, SACT dataset
Radiotherapy treatment	Derived based on any record of radiotherapy (excluding brachytherapy (rttreatmentmodality=06)) Captured rtprescribeddose & prescribedfractions to flag palliative regimens of 30Gy/15 fractions; 26Gy/5 fractions; 20Gy/5 fractions; 8Gy/1 fractions	RTDS	CaNISC: Derived based on the presence of a value in the variable START_DATE_OF_RADIOOTHERAPY RT dataset: Derived based on the presence of a value in the variable RTStartDate Nb: RT dataset provided for people diagnosed in the 2023 calendar year only For RT fractionation information: - From CaNISC: RADIOOTHERAPY_PRESCRIBED_DOSE and RADIOOTHERAPY_PRESCRIBED_FRACTION - From RT dataset: PrescribedDoseTotal and PrescribedFractionTotal	CaNISC, RT dataset
Radiotherapy date	apptdate associated with radiotherapy treatment	RTDS	CaNISC: START_DATE_OF_RADIOOTHERAPY RT dataset:	CaNISC, RT dataset
Chemoradiotherapy treatment	Derived based on record of SACT and radiotherapy	Derived	Derived based on record of SACT and radiotherapy	Derived
Disease-targeted treatment	Any record of surgery, SACT, or radiotherapy up to 9 months (274 days) after diagnosis date Note: records of surgery up to 30 days before diagnosis date were also permitted	Derived	Any record of surgery, SACT, or radiotherapy up to 9 months (274 days) after diagnosis date Note: records of surgery up to 30 days before diagnosis date were also permitted	Derived
First treatment date	Earliest of surgery date, SACT date, and radiotherapy date up to 9 months (274 days) after diagnosis date Note: records of surgery up to 30 days before diagnosis date were also permitted	Derived	Earliest of surgery date, SACT date, and radiotherapy date up to 9 months (274 days) after diagnosis date Note: records of surgery up to 30 days before diagnosis date were also permitted	Derived
<i>Supportive care for pancreatic cancer</i>				
CNS (Clinical Nurse Specialist) involved	Derived using clinicalnursespecialist, counting any “Yes” response option as CNS involved	Derived by NDRS from COSD	Data not provided	
PERT (Pancreatic Enzyme Replacement	Derived based on prescribedbnfcode=0109040 or prescribedbnfname of	Primary care	Data not provided	

Treatment prescription)	"Creon", "Pancrease", "Nutrizym", or "Pancrex"	prescribing data		
<i>Survival outcomes</i>				
Date of death	Derived based on vitalstatusdate when vitalstatus=D If vitalstatusdate missing, replace with deathdatebest if available Note: some people will not have a date of death if at the last known vitalstatusdate their vitalstatus=A (Alive)	NCRD	Date of death derived from the earliest date of two variables: DATE_OF_DEATH (CaNISC) and date_of_death (ONS)	CaNISC , ONS
Survival at specific time points post-diagnosis	Derived based on difference between diagnosis date and vital status date Example: 30 days post diagnosis, person flagged as deceased if vitalstatusdate minus diagnosis date <30 & vitalstatus=D (dead); person flagged as alive if vitalstatusdate minus diagnosis date >30		Calculated using time between DIAGNOSIS_DATE (CaNISC) and the date of death. Date of death derived from the earliest date of two variables: DATE_OF_DEATH (CaNISC) and date_of_death (ONS)	CaNISC , ONS

5. Indicator Definitions

The audit uses key indicators to monitor progress against its healthcare improvement goals. These indicators align with national guidelines and standards. Definitions of how the indicators included in the SotN report were derived from data for England and Wales are described below.

Indicators were reported for people with a pancreatic cancer diagnosis code. A sub-set of indicators were reported for people with an extrahepatic bile duct or ampullary tumour.

5.1 Performance Indicator 1: FDG-PET/CT scan prior to surgery

Percentage of people who had an FDG/PET-CT scan prior to surgery for pancreatic cancer.

Table 5.1: Percentage of people who had an FDG-PET/CT scan prior to surgery for pancreatic cancer		
	<u>England</u>	<u>Wales</u>
Dates of diagnosis:	1/1/2020 to 31/12/2022	1/1/2022 to 31/12/2023
Numerator:	Number of people with an imaging record for an FDG-PET/CT scan up to 6 months prior to surgery record	Number of people with a record of a date of an FDG-PET/CT (Imaging__PET) in the 6 months prior to any pancreatic

Number of people with a record of an FDG-PET/CT prior to surgery		surgery occurring in a Welsh local health board.
Denominator: Number of people with a record of any pancreatic surgery	Number of people with a record of any pancreatic surgery within 30 days prior to diagnosis date or up to 9 months after diagnosis date	Number of people with a record of any pancreatic surgery occurring in a Welsh local health board within 30 days prior to diagnosis date or up to 9 months after diagnosis date
Construction notes	Time restriction: only count imaging records on or before date of pancreatic surgery, and no more than 183 days (6 months) prior to date of pancreatic surgery Liver MRI reported separately as additional information, but not included within the indicator	Time restriction: only count imaging records on or before date of pancreatic surgery, and no more than 183 days (6 months) prior to date of any pancreatic surgery. Details on the type of PET scan performed were not provided (eg. tracer used or anatomical site scanned). We have therefore included anyone with a record of any PET scan in this indicator.
Country reporting:	England & Wales separately	
Organisational Reporting level:	HPB specialist centre	National
Subgroup Reporting:	None	None
Risk adjusted:	No	
Outlier reporting:	No	

5.2 Performance Indicator 2: Multidisciplinary team (MDT) meeting

Percentage of people who had a record of being discussed at an MDT (Multidisciplinary Team) meeting.

Table 5.2: Percentage of people who had a record of being discussed at a multidisciplinary team (MDT) meeting		
	<u>England</u>	<u>Wales</u>
Dates of diagnosis:	1/1/2021 to 31/12/2022	Not reported
Numerator:	Number of people with a record of an MDT meeting date within 60 days before or after diagnosis date	Not reported
Denominator:	Number of people with a primary diagnosis of pancreatic cancer	Not reported
Construction notes		
Country reporting:	England only	
Organisational Reporting level:	Trust of diagnosis Cancer Alliance of diagnosis	Not reported
Subgroup Reporting:	None	Not reported
Risk adjusted:	No	Not reported
Outlier reporting:	No	Not reported

5.3 Performance Indicator 3: Biliary stent prior to surgery

Percentage of people undergoing a Whipple procedure (without neoadjuvant chemotherapy) who had a biliary stent placed prior to surgery.

Table 5.3: Percentage of people undergoing a Whipple procedure (without neoadjuvant chemotherapy) who had a biliary stent placed prior to surgery

	<u>England</u>	<u>Wales</u>
Dates of diagnosis:	1/1/2020 to 31/12/2022	1/1/2022 to 31/12/2023
Numerator: Number of people with a record of biliary stent prior to Whipple procedure without record of SACT/RT before procedure	Number of people with a record of biliary stent prior to Whipple procedure without record of SACT/RT up to 14 weeks (98 days) before procedure	Number of people with a record of biliary stent prior to Whipple procedure in a Welsh health board without record of SACT/RT up to 14 weeks (98 days) before procedure
Denominator: Number of people who had a Whipple procedure without a record of SACT/RT before procedure	Number of people who had a Whipple procedure without a record of SACT/RT up to 14 weeks (98 days) before procedure	Number of people who had a Whipple procedure in a Welsh health board only, without a record of SACT/RT up to 14 weeks (98 days) before procedure
Construction notes	Neo-adjuvant chemo-radiotherapy: see approach as part of indicator #7	Neo-adjuvant chemo-radiotherapy: see approach as part of indicator #7
Country reporting:	England & Wales separately	
Organisational Reporting level:	HPB specialist centre	National
Subgroup Reporting:	None	None
Risk adjusted:	No	
Outlier reporting:	No	

5.4 Performance Indicator 4: Time to treatment

Time from urgent suspected cancer GP referral to first disease-targeted treatment, presented as:

- Time from GP referral to diagnosis
- Time from GP referral to first treatment
- Time from diagnosis to first treatment (for those referred via GP)
- % diagnosed within 21 and 28 days of GP referral

Table 5.4: Time from referral to first treatment (days)

	<u>England</u>	<u>Wales</u>
Dates of diagnosis:	1/1/2021 to 31/12/2022	1/1/2022 to 31/12/2023
Numerator: Median time (days) from urgent suspected cancer GP referral to first treatment	Median time (days) from urgent GP referral to first disease-targeted treatment	Median time (days) from urgent suspected cancer GP referral to first disease-targeted treatment
Denominator: N/A	N/A	N/A
Construction notes	Calculate median and IQR for the following time periods: • Urgent GP referral date to diagnosis date	Calculate median and IQR for the following time periods: • GP referral date to diagnosis date • GP referral date to date of first disease-targeted treatment date

	- Urgent GP referral date to date of first disease-targeted treatment date - Diagnosis date to first disease-targeted treatment date (for those with an urgent GP referral) Calculate % with a diagnosis date within 21 and 28 days or urgent GP referral date: Numerator: count of people with ≤21 (28) days between date of urgent GP referral and diagnosis date Denominator: count of people with an urgent GP referral date	Diagnosis date to first disease-targeted treatment date (for those with an GP referral) Source of referral: ("Referral from General Medical Practitioner" in the variable SOURCE_OF_REFERRAL (CaNISC) Calculate % with a diagnosis date within 21 and 28 days or urgent GP referral date: Numerator: count of people with ≤21 (28) days between date of GP referral and diagnosis date Denominator: count of people with an GP referral date
Country reporting:	England & Wales Combined	
Organisational Reporting level:	Trust of diagnosis Cancer Alliance of diagnosis	Health board of diagnosis
Subgroup Reporting:	None	None
Risk adjusted:	No	
Outlier reporting:	No	

5.5 Performance Indicator 5: Receipt of disease-targeted treatment (stage 1-3)

Percentage of people with non-metastatic (stage 1-3) pancreatic cancer who receive disease-targeted treatment (surgery, SACT, radiotherapy).

Table 5.5: Percentage of people with non-metastatic (stage 1-3) pancreatic cancer who received disease-targeted treatment

	<u>England</u>	<u>Wales</u>
Dates of diagnosis:	1/1/2021 to 31/12/2022	1/1/2022 to 31/12/2023
Numerator: Number of people with a record of disease-targeted treatment	Number of people with a record of disease-targeted treatment	Number of people with a record of disease-targeted treatment
Denominator: Number of people diagnosed with stage 1-3 (non-metastatic) pancreatic cancer	Number of people diagnosed with stage 1-3 (non-metastatic) pancreatic cancer	Number of people diagnosed with stage 1-3 or M0 pancreatic cancer
Construction notes		Surgeries of patients diagnosed in Wales were included regardless of whether the surgery took place in England or Wales.
Country reporting:	England & Wales separately	
Organisational Reporting level:	Trust of diagnosis Cancer Alliance of diagnosis	Health board of diagnosis
Subgroup Reporting:	None	None
Risk adjusted:	No	
Outlier reporting:	No	

5.6 Performance Indicator 6: Receipt of disease-targeted treatment (stage 4)

Percentage of people with metastatic (stage 4) pancreatic cancer who receive disease-targeted treatment (surgery, SACT, radiotherapy).

Table 5.6: Percentage of people with metastatic (stage 4) pancreatic cancer who received disease-targeted treatment		
	<u>England</u>	<u>Wales</u>
Dates of diagnosis:	1/1/2021 to 31/12/2022	1/1/2022 to 31/12/2023
Numerator: Number of people with a record of disease-targeted treatment	Number of people with a record of disease-targeted treatment	Number of people with a record of disease-targeted treatment
Denominator: Number of people diagnosed with stage 4 (metastatic) pancreatic cancer	Number of people diagnosed with stage 4 (metastatic) pancreatic cancer	Number of people diagnosed with stage 4 or M1 pancreatic cancer
Construction notes		Surgeries of patients diagnosed in Wales were included regardless of whether the surgery took place in England or Wales.
Country reporting:	England & Wales separately	
Organisational Reporting level:	Trust of diagnosis Cancer Alliance of diagnosis	Health board of diagnosis
Subgroup Reporting:	None	None
Risk adjusted:	No	
Outlier reporting:	No	

5.7 Performance Indicator 7: Receipt of chemotherapy and/or radiotherapy alongside surgery

Percentage of people with pancreatic cancer receiving SACT/RT alongside surgery.

Table 5.7: Percentage of people with pancreatic cancer who received chemotherapy and/or radiotherapy alongside surgery		
	<u>England</u>	<u>Wales</u>
Dates of diagnosis:	1/1/2020 to 31/12/2022	1/1/2022 to 31/12/2023
Numerator: Number of people who received SACT and/or radiotherapy (a) before surgery or (b) after Whipple procedure	a. Before surgery: number of people who received SACT and/or radiotherapy up to 14 weeks before any pancreatic surgery b. After surgery: Number of people who received SACT and/or radiotherapy up to 14 weeks after Whipple procedure	a. Before surgery: number of people who received SACT and/or radiotherapy up to 14 weeks (98 days) before any pancreatic surgery b. After surgery: Number of people who received SACT and/or radiotherapy up to 14 weeks (98 days) after Whipple procedure

Denominator: Number of people who underwent (a) any pancreatic surgery or (b) Whipple procedure	a. Before surgery: number of people with a record of any pancreatic surgery 30 days before or 9 months after diagnosis date b. After surgery: number of people with a record of a Whipple procedure 30 days before or 9 months after diagnosis date	a. Before surgery: number of people who underwent any pancreatic surgery 30 days before or 9 months after diagnosis date in England or Wales b. After surgery: number of people who underwent a Whipple procedure 30 days before or 9 months after diagnosis date in England or Wales
Construction notes	SACT/RT before surgery: count when at least one of the following is recorded: (1) any SACT treatment when SACT date <=98 days before pancreatic surgery date, (2) any radiotherapy treatment when radiotherapy date <=98 days before pancreatic surgery date SACT/RT after Whipple: count when at least one of the following is recorded: (1) any SACT treatment when SACT date <= 98 days after Whipple procedure date, (2) any radiotherapy treatment, excluding palliative doses*, when radiotherapy date <=98 days after Whipple procedure date * Following combinations of rtprescribeddose & prescribedfractions considered palliative: 30Gy/15 fractions; 26Gy/5 fractions; 20Gy/5 fractions; 8Gy/1 fractions	• SACT/RT before surgery: count when at least one of the following is recorded ○ Any SACT treatment when SACT date <=98 days before pancreatic surgery date ○ Any radiotherapy treatment when radiotherapy date <=98 days before pancreatic surgery date • SACT/RT after Whipple: count when at least one of the following is recorded: ○ Any SACT treatment when SACT date <= 98 days after Whipple procedure date ○ Any radiotherapy treatment, excluding palliative doses**, when radiotherapy date <=98 days after Whipple procedure date The following combinations of radiotherapy were considered palliative: 30Gy/15 fractions; 26Gy/5 fractions; 20Gy/5 fractions; 8Gy/1 fractions
Country reporting:	England & Wales separately	
Organisational Reporting level:	HPB specialist centre	Health board of diagnosis
Subgroup Reporting:	None	None
Risk adjusted:	No	
Outlier reporting:	No	

5.8 Performance Indicator 8: Clinical nurse specialist (CNS) involvement

Percentage of people with pancreatic cancer who were seen by a clinical nurse specialist (CNS).

Table 5.8: Percentage of people with pancreatic cancer who were seen by a clinical nurse specialist (CNS)

	<u>England</u>	<u>Wales</u>
--	----------------	--------------

Dates of diagnosis:	1/1/2021 to 31/12/2022	Not reported
Numerator: Number of people with CNS involved	Number of people with CNS involved	Not reported
Denominator: Number of people with a primary diagnosis of pancreatic cancer with complete information related to CNS	Number of people with a primary diagnosis of pancreatic cancer with complete information related to CNS	Not reported
Construction notes:		
Country reporting:	England only	
Organisational Reporting level:	Trust of diagnosis Cancer Alliance of diagnosis	Not reported
Subgroup Reporting:	None	Not reported
Risk adjusted:	No	Not reported
Outlier reporting:	No	Not reported

5.9 Performance Indicator 9: Pancreatic enzyme replacement therapy (PERT) use

Percentage of people who were prescribed pancreatic enzyme replacement therapy (PERT) in primary care.

Table 5.9: Percentage of people who were prescribed pancreatic enzyme replacement therapy (PERT) in primary care		
	<u>England</u>	<u>Wales</u>
Dates of diagnosis:	1/1/2021 to 31/12/2022	Not reported
Numerator: Number of people with a prescription of PERT	Number of people with a prescription of PERT in primary care	Not reported
Denominator: Number of people with a primary diagnosis of pancreatic cancer	Number of people with a primary diagnosis of pancreatic cancer	Not reported
Construction notes:		
Country reporting:	England only	
Organisational Reporting level:	Trust of diagnosis Cancer Alliance of diagnosis	Not reported
Subgroup Reporting:	None	Not reported
Risk adjusted:	No	Not reported
Outlier reporting:	No	Not reported

5.10 Performance Indicator 10: Survival from diagnosis

Survival at 30- and 90 days, and 1- and 2 years after diagnosis.

Table 5.10: Survival at 30- and 90-days, and 1- and 2-years after diagnosis		
	<u>England</u>	<u>Wales</u>

Dates of diagnosis:	1/1/2021 to 31/12/2022 (for 30-, 90-day, and 1-year) 1/1/2020 to 31/12/2021 (for 2-year)	1/1/2022 to 31/12/2023
Numerator: Number of people alive at specific time points after diagnosis of pancreatic cancer	Number of people alive at 30 days, 90 days, 1 year, and 2 years after diagnosis of pancreatic cancer	Number of people alive more than 30 days, 90 days and 1 year after diagnosis of pancreatic cancer
Denominator: Number of people with a primary diagnosis of pancreatic cancer and a vital status date	Number of people with a primary diagnosis of pancreatic cancer and a vital status date	For overall survival: number of people with diagnosis of pancreatic cancer
Construction notes:		
Country reporting:	England & Wales separately	
Organisational Reporting level:	Trust of diagnosis Cancer Alliance of diagnosis	Health board of diagnosis
Subgroup Reporting:	By stage at diagnosis	By stage at diagnosis
Risk adjusted:	Yes: age group at diagnosis, sex, index of multiple deprivation quintile, stage at diagnosis, performance status at diagnosis, receipt of disease targeted treatment, RCS Charlson Comorbidity Index, and diagnosis year	
Outlier reporting:	Yes: 90-day and 1-year survival indicators	

6. NHS organisations

The audit presents organisation-level findings by the NHS organisation of diagnosis or treatment, as appropriate for the specific indicator:

- Organisation of surgery: for indicators concerned with surgery
- Organisation of diagnosis: for all other indicators (in England this includes trust of diagnosis and Cancer Alliance of diagnosis)

Details on allocation to NHS Trust in England:

- Organisation of surgery are the HPB specialist centres identified via the organisation codes listed in Appendix 3: Table 7
- Trust of diagnosis is identified using the trust of diagnosis variable in the NCRD.
- In instances where the trust of diagnosis is a tertiary centre (The Christie, The Clatterbridge, or The Royal Marsden), these cases are not reported at the trust-level for The Christie or The Clatterbridge (as they are not diagnosing trusts) but they are included in Cancer Alliance and National-level reporting.
- Cases where the organisation of diagnosis is not an NHS organisation are excluded from the Trust and Cancer Alliance-level analyses but are included in national-level results.
- Cases in the English data where the organisation of diagnosis is an NHS Wales organisation are excluded from the England analyses, unless the case had surgery in England.

For Wales, the local health board of diagnosis was identified using the organisation codes listed in Table 9.

A minimum of five diagnoses in the audit period were required for reporting at trust or health board level. This was to ensure only trusts providing cancer services were included and also to avoid very small numbers which can lead to unreliable estimates and increase the risk of potential data disclosure.

7. Statistical Analysis

All statistical analyses were conducted using STATA version 17.0.

Most results in the SotN Report are descriptive. The results of categorical data items are reported as percentages (%). Results are typically provided as an overall figure with an indication of variation by organisational reporting level (see NHS organisations section). Note that within tables in the SotN Report, the total percentage may not equal 100%, due to rounding.

7.1 Small number Suppression

- Data quality and completeness results have not been suppressed.
- For results presented at organisational level, cell values are suppressed when there are less than 25 diagnoses at the organisation and/or the denominator was <10.

7.2 Risk-adjustment of indicators

The tables of performance indicators state whether risk adjustment has been performed.

Multivariable logistic regression models were used to estimate the likelihood of survival for each individual who had a diagnosis of pancreatic cancer (based on their characteristics), and these probabilities have been summed to calculate the predicted number of people surviving for each organisation. The regression models include the following patient characteristics: age group at diagnosis, sex, index of multiple deprivation, stage at diagnosis, performance status at diagnosis, receipt of disease targeted treatment, RCS Charlson Comorbidity Index, and diagnosis year. Data for England and Wales were analysed separately.

Risk adjusted rates are presented only for organisations with at least 10 people diagnosed during the relevant period.

Table 7.1 below provides details on the datasets and variables used in the risk adjustment model. See section **Error! Reference source not found.** for further details on construction notes for any of the variables listed.

Table 7.1: Risk Adjustment Variables		
Data Item	Additional detail	
	England	Wales
Age at diagnosis	Categorised into age groups: <60 years, 60-69 years, 70-79 years, 80+ years	Categorised into age groups: <60 years, 60-69 years, 70-79 years, 80+ years
Sex	No additional detail	No additional detail
Index of multiple deprivation	No additional detail	No additional detail
Stage at diagnosis	No additional detail	No additional detail
Performance status at diagnosis	Categorised into the following groups: 0, 1, 2, 3/4	No additional detail
Receipt of disease targeted treatment	Binary variable, yes/no	Binary variable, yes/no

Charlson comorbidity index	The CCI is a commonly used scoring system for medical comorbidities, consisting of a grouped score calculated based on the absence (0) and presence (≥ 1) of 14 pre-specified medical conditions (Appendix 2). The CCI was calculated using information on secondary diagnoses (ICD-10 codes) recorded in HES APC (England) / PEDW (Wales) within the 12-month period prior to a patient's diagnosis. For the purpose of analysis, the CCI is grouped into four categories: • 0 none of the 14 pre-specified comorbidities. • 1 only 1 of the 14 pre-specified comorbidities. • 2 only 2 of the 14 pre-specified comorbidities • 3+ 3 or more of the 14 pre-specified comorbidities In instances where there were no HES-APC records for a person included in the audit, the variable <code>chrl_tot_27_03</code> from NCRD was used as the CCI value.	
Diagnosis year	Year extracted from diagnosis date	Year extracted from DIAGNOSIS_DATE (CaNISC)

7.3 Handling of missing data

For the risk-adjustment, missing values for stage, performance status, IMD quintile (Wales only) and age (Wales only) were imputed with multiple imputation using chained equations, creating ten data sets and pooling model estimates using Rubin's Rules. The imputation models included all the variables in the analysis models.

8. Outlier Process

The outlier process can be found in the separate [audit outlier policy](#).

Appendix 1: Routine data sources

Overview of the data sources used for the SotN Report.

Country	Data source	Content
England	Cancer registry (NCRD)	Data on all aspects of the cancer registration including information from hospital pathology systems.
England	COSD	Cancer Outcomes and Services dataset (COSD) items are submitted routinely by service providers via multidisciplinary team (MDT) electronic data collection systems to the National Cancer Data Repository (NCDR) on a monthly basis.
England	SACT	Systemic Anti-Cancer Therapy (SACT) data contains information on chemotherapy dates, regimen(s) and dose(s).
England	RTDS	Radiotherapy dataset (RTDS) contains information on radiotherapy treatment including dates, prescription region and dose.
England	HES	Hospital Episode Statistics (HES) is the administrative database of all NHS hospital admissions in England. The audit receives records from both admitted patient care (HES-APC) and outpatient care (HES-OP).
England	PCPD	Primary Care Prescription Database (PCPD) contains information on prescriptions in primary care.
England	CWT	Cancer Waiting Times (CWT) contains information on dates and sources of referrals, diagnoses, and treatments for cancer. This information is uploaded monthly by NHS providers and is used to monitor cancer waiting times.
England	DIDs	The Diagnostic Imaging Dataset (DID) contains detailed information about diagnostic imaging tests carried out for NHS patients, including details of the test (type of test and body site) and date of imaging. Information is extracted from local radiology information systems.
Wales	CaNISC	Cancer Network Information System Cymru (Canisc) contains data on all aspects of the cancer registration including investigations.
Wales	PEDW	Patient Episode Database for Wales (PEDW) is the administrative database of all NHS hospital admissions in Wales. Nb: not every person in the cancer registry had a linked PEDW record.
Wales	LSOA	Lower-layer Super Output Areas (LSOA) dataset contains information on deprivation in small areas (LSOAs) across Wales.
Wales	RTH	Radiotherapy data (RTH) contains information on radiotherapy treatment.
Wales	SACT	SACT data contains information on systemic anti-cancer treatment.
Wales	ONS	Office for National Statistics (ONS) death data including date of death and cause of death.

Appendix 2: Charlson Comorbidity Index

Reference:

Armitage JN, van der Meulen JH. Identifying co-morbidity in surgical patients using administrative data with the Royal College of Surgeons Charlson Score. *Br J Surg* 2010;97:772-81. doi <https://doi.org/10.1002/bjs.6930>

Pre-specified conditions included in the assignment of Charlson Comorbidity Index (CCI).

CCI Conditions
Myocardial infarction
Dementia
Diabetes mellitus
Metastatic solid tumour
Congestive cardiac failure
Chronic pulmonary disease
Hemiplegia or paraplegia
AIDS/HIV infection
Peripheral vascular disease
Rheumatological disease
Renal disease
Cerebrovascular disease
Liver disease
Any malignancy

Note: AIDS/HIV diagnoses cannot be identified in HES APC data because of legal requirements for NHS trusts to remove patient identifiers from legally restricted records, including those containing diagnoses of HIV/AIDS. These diagnoses are also not found in linked PEDW data.

Appendix 3: Code lists

Table 1. ICD-10 codes used to define pancreatic cancer audit cohort

Code	Description
<i>Malignant neoplasm of pancreas</i>	
C25.0	Head of pancreas
C25.1	Body of pancreas
C25.2	Tail of pancreas
C25.3	Pancreatic duct
C25.7	Other parts of pancreas: neck of pancreas
C25.8	Overlapping lesion of pancreas
C25.9	Pancreas, unspecified
<i>Malignant neoplasm of extrahepatic bile duct and ampullary tumours</i>	
C24.0	Extrahepatic bile duct: biliary duct or passage NOS, common bile duct, cystic duct, hepatic duct
C24.1	Ampulla of Vater

Source of ICD-10 codes: <https://icd.who.int/browse10/2019>

Table 2. Morphology codes for identification of neuroendocrine tumours

Code	Description
8013	Large cell neuroendocrine carcinoma
8041	Small cell carcinoma, NOS
8042	Oat cell carcinoma
8043	Small cell carcinoma, fusiform cell
8044	Small cell carcinoma, intermediate cell
8045	Combined small cell carcinoma
8150	Islet cell carcinoma
8151	Insulinoma
8152	Glucagonoma
8153	Gastrinoma
8154	Mixed islet cell & exocrine adenocarcinoma
8155	Vipoma
8156	Somatostatinoma
8157	Enteroglucagonoma
8158	ACTH-producing tumor
8240	Carcinoid tumour
8241	Enterochromaffin cell carcinoid
8242	Enterochromaffin-like cell tumour
8243	Goblet cell carcinoid
8244	Composite carcinoid
8245	Adenocarcinoid tumour
8246	Neuroendocrine carcinoma
8247	Merkel cell carcinoma
8249	Atypical carcinoid tumour
9091	Strumal carcinoid

Source of morphology codes: <https://biobank.ndph.ox.ac.uk/ukb/ukb/docs/ICDcancermorph.pdf>

Reference publication - neuroendocrine morphology codes: <https://www.nature.com/articles/s41416-019-0606-3>

Table 3. SNOMED codes used to identify scans

SNOMED-CT ID	Description
<i>FDG-PET/CT</i>	
725928006	Positron emission tomography with computed tomography fluorodeoxyglucose F18 imaging of base of brain to mid-thigh (procedure)
443271005	Positron emission tomography with computed tomography using fluorodeoxyglucose (18-F) (procedure)
432675001	Positron emission tomography fluorodeoxyglucose imaging of whole body (procedure)
1097781000000108	Positron emission tomography with computed tomography 18F fluorodeoxyglucose imaging of cranial vertex to mid-thigh (procedure)
<i>Liver MRI</i>	
910561000000103	Diffusion weighted magnetic resonance imaging of liver (procedure)
432551009	Magnetic resonance imaging of liver and spleen (procedure)
431839003	Magnetic resonance imaging of liver with contrast (procedure)
432633002	Magnetic resonance imaging of liver and biliary tract with contrast (procedure)
764569004	Magnetic resonance imaging of liver and spleen with contrast (procedure)
1065681000000100	Magnetic resonance imaging of liver and spleen with contrast (procedure)
911811000000107	Magnetic resonance imaging of transplanted liver (procedure)
241622002	Magnetic resonance imaging of liver (procedure)

Source of SNOMED-CT codes for diagnostic imaging (Annex 5): <https://www.england.nhs.uk/statistics/statistical-work-areas/diagnostic-imaging-dataset/>

Table 4. OPCS-4 codes used to identify pancreatic surgery

OPCS-4 code	Description
<i>Whipple procedure</i>	
J56.1	Pancreaticoduodenectomy and excision of surrounding tissue
J56.2	Pancreaticoduodenectomy and resection of antrum of stomach
J56.3	Pancreaticoduodenectomy NEC
J56.4	Subtotal excision of head of pancreas with preservation of duodenum and drainage HFQ
<i>All other pancreatic surgeries</i>	
J55.1	Total pancreatectomy and excision of surrounding tissue
J55.2	Total pancreatectomy NEC
J55.8	Total excision of pancreas, other specified
J55.9	Total excision of pancreas, unspecified
J56.8	Excision of head of pancreas, other specified
J56.9	Excision of head of pancreas, unspecified
J57.1	Subtotal pancreatectomy
J57.2	Left pancreatectomy and drainage of pancreatic duct
J57.3	Left pancreatectomy NEC
J57.4	Excision of tail of pancreas and drainage of pancreatic duct
J57.5	Excision of tail of pancreas NEC
J57.8	Other partial excision of pancreas, other specified
J57.9	Other partial excision of pancreas, unspecified

Source of OPCS-4 codes: <https://classbrowser.nhs.uk/#/book/OPCS-4.10/>

Table 5. OPCS-4 codes used to identify biliary stents

OPCS-4 code	Description
J382	Endoscopic sphincterotomy of sphincter of Oddi and insertion of tubal prosthesis into bile duct
J394	Endoscopic sphincteroplasty of ampulla of Vater and insertion of tubal prosthesis into bile duct
J401	Endoscopic retrograde insertion of tubal prosthesis into both hepatic ducts
J402	Endoscopic retrograde insertion of tubal prosthesis into bile duct NEC
J403	Endoscopic retrograde renewal of tubal prosthesis in bile duct NEC
J405	Endoscopic retrograde insertion of expanding covered metal stent into bile duct
J406	Endoscopic retrograde insertion of expanding metal stent into bile duct NEC
J407	Endoscopic retrograde renewal of expanding metal stent in bile duct
J408	Other specified endoscopic retrograde placement of prosthesis in bile duct
J409	Unspecified endoscopic retrograde placement of prosthesis in bile duct
J418	Other specified other therapeutic endoscopic retrograde operations on bile duct
J431	Endoscopic retrograde cholangiopancreatography and biopsy of lesion of ampulla of Vater
J432	Endoscopic retrograde cholangiopancreatography and biopsy of lesion of biliary or pancreatic system NEC
J433	Endoscopic retrograde cholangiopancreatography and collection of bile
J438	Other specified diagnostic endoscopic retrograde examination of bile duct and pancreatic duct
J439	Unspecified diagnostic endoscopic retrograde examination of bile duct and pancreatic duct
J441	Endoscopic retrograde cholangiography and biopsy of lesion of bile duct
J448	Other specified diagnostic endoscopic retrograde examination of bile duct
J449	Unspecified diagnostic endoscopic retrograde examination of bile duct
J471	Percutaneous insertion of tubal prosthesis into both hepatic ducts
J472	Percutaneous insertion of tubal prosthesis into right hepatic duct NEC
J473	Percutaneous insertion of tubal prosthesis into left hepatic duct NEC
J474	Percutaneous insertion of tubal prosthesis into hepatic duct NEC
J475	Percutaneous insertion of tubal prosthesis into common bile duct
J478	Other specified therapeutic percutaneous insertion of prosthesis into bile duct
J479	Unspecified therapeutic percutaneous insertion of prosthesis into bile duct
J481	Renewal of percutaneously inserted tubal prosthesis in bile duct
J483	Attention to percutaneously inserted tubal prosthesis in bile duct NEC
J485	Percutaneous transhepatic biliary drainage multiple
J486	Percutaneous transhepatic biliary drainage single
J488	Other specified other therapeutic percutaneous operations on bile duct
J489	Unspecified other therapeutic percutaneous operations on bile duct
J502	Percutaneous cholangiography NEC
J505	Percutaneous transhepatic cholangiography

Source of OPCS-4 codes: <https://classbrowser.nhs.uk/#/book/OPCS-4.10/>

Table 6. Excluded regimens in Systemic Anti-Cancer Therapy (SACT) dataset

Regimens excluded from SACT analyses
benchmark_group= "NOT CHEMO" & none in drug_group are systemic anti-cancer treatments
benchmark_group= "LUTETIUM-177"
benchmark_group= "ZOLEDRONIC ACID" & none in drug_group are systemic anti-cancer treatments

benchmark_group= "DENOSUMAB" & none in drug_group are systemic anti-cancer treatments
benchmark_group= "HORMONES" & drug_group contains "hormones"
benchmark_group= "PAMIDRONATE" & no other drugs listed in drug_group
benchmark_group= "STREPTOZOCIN" & none in drug_group are systemic anti-cancer treatments

Table 7. Trust codes for HPB specialist centres

Trust code	Trust name
RTD	The Newcastle Upon Tyne Hospitals NHS Foundation Trust
REM	Liverpool University Hospitals NHS Foundation Trust
RJE	University Hospitals of North Midlands NHS Trust
RKB	University Hospitals Coventry and Warwickshire NHS Trust
RRK	University Hospitals Birmingham NHS Foundation Trust
RJZ	King's College Hospital NHS Foundation Trust
RA2	Royal Surrey County Hospital NHS Foundation Trust
RTH	Oxford University Hospitals NHS Foundation Trust
RK9	University Hospitals Plymouth NHS Trust
RA7	University Hospitals Bristol and Weston NHS Foundation Trust
RHM	University Hospital Southampton NHS Foundation Trust
RXR	East Lancashire Hospitals NHS Trust
ROA	Manchester University NHS Foundation Trust
RPY	The Royal Marsden NHS Foundation Trust
RYJ	Imperial College Healthcare NHS Trust
RGT	Cambridge University Hospitals NHS Foundation Trust
RWE	University Hospitals of Leicester NHS Trust
RX1	Nottingham University Hospitals NHS Trust
RHQ	Sheffield Teaching Hospitals NHS Foundation Trust
RWA	Hull University teaching Hospitals NHS Trust
RAL	Royal Free London NHS Foundation Trust
R1H	Barts Health NHS Trust
RR8	Leeds Teaching Hospitals NHS Trust

Source: <https://www.pancreaticcancer.org.uk/support-for-you/your-care/your-local-pancreatic-cancer-specialist-centre/>

Table 8. Codes used to identify SACT in HES and PEDW (OPCS-4 codes)

OPCS-4 code	Description
X701	Procurement of drugs for chemotherapy for neoplasm for regimens in Band 1
X702	Procurement of drugs for chemotherapy for neoplasm for regimens in Band 2
X703	Procurement of drugs for chemotherapy for neoplasm for regimens in Band 3
X704	Procurement of drugs for chemotherapy for neoplasm for regimens in Band 4
X705	Procurement of drugs for chemotherapy for neoplasm for regimens in Band 5
X708	Procurement of drugs for chemotherapy for neoplasm in Bands 1-5: Other specified
X709	Procurement of drugs for chemotherapy for neoplasm in Bands 1-5: Unspecified
X711	Procurement of drugs for chemotherapy for neoplasm for regimens in Band 6
X712	Procurement of drugs for chemotherapy for neoplasm for regimens in Band 7
X713	Procurement of drugs for chemotherapy for neoplasm for regimens in Band 8
X714	Procurement of drugs for chemotherapy for neoplasm for regimens in Band 9
X715	Procurement of drugs for chemotherapy for neoplasm for regimens in Band 10
X718	Procurement of drugs for chemotherapy for neoplasm in Bands 6-10: Other specified
X719	Procurement of drugs for chemotherapy for neoplasm in Bands 6-10: Unspecified

X721	Delivery of complex chemotherapy for neoplasm including prolonged infusional treatment at first attendance
X722	Delivery of complex parenteral chemotherapy for neoplasm at first attendance
X723	Delivery of simple parenteral chemotherapy for neoplasm at first attendance
X724	Delivery of subsequent element of cycle of chemotherapy for neoplasm
X728	Delivery of chemotherapy for neoplasm: Other specified
X729	Delivery of chemotherapy for neoplasm: Unspecified
X731	Delivery of exclusively oral chemotherapy for neoplasm
X738	Delivery of oral chemotherapy for neoplasm: Other specified
X739	Delivery of oral chemotherapy for neoplasm: Unspecified
X748	Other chemotherapy drugs: Other specified
X749	Other chemotherapy drugs: Unspecified
X352	Intravenous chemotherapy
X353	Intravenous immunotherapy
X373	Intramuscular chemotherapy
X374	Intramuscular immunotherapy
X384	Subcutaneous chemotherapy
X385	Subcutaneous immunotherapy

Table 9. Organisation codes for Wales Health Boards

Organisation code	Hospital code	Trust name
7A1	7A1A1	Glan Clwyd Hospital
7A1	7A1A4	Wrexham Maelor Hospital
7A1	7A1AU	Ysbyty Gwynedd
7A2	7A2AG	Glangwili General Hospital
7A2	7A2AJ	Bronglais General Hospital
7A2	7A2AL	Prince Philip Hospital
7A2	7A2BL	Withybush General Hospital
7A3	7A3C4	Singleton Hospital
7A3	7A3C7	Morriston Hospital
7A3	7A3CJ	Neath Port Talbot Hospital
7A4	7A4BV	University Hospital of Wales
7A4	7A4C1	University Hospital Llandough
7A5	7A3B7	Princess of Wales Hospital
7A5	7A5B1	Royal Glamorgan Hospital
7A5	7A5B3	Prince Charles Hospital
7A6	7A6AM	Nevill Hall Hospital
7A6	7A6AR	Royal Gwent Hospital
7A6	7A6G9	The Grange University Hospital
RQF		VELINDRE NHS TRUST

Appendix 4: AJCC TNM staging of (exocrine) pancreatic cancer

Stage	T	N	M
1	1	0	0
	2	0	0
2	3	0	0
	1-3	1	0
3	4	0-2	0
	1-4	2	0
4	1-4	0-2	1

Reference: MB Amin, SB Edge, FL Greene, et al, eds. AJCC Cancer Staging Manual. 8th ed. New York: Springer; 2017

Appendix 5: World Health Organisation Performance Status

Performance status	Definition
0	Able to carry out all normal activity without restriction
1	Restricted in strenuous activity but ambulatory and able to carry out light work
2	Ambulatory and capable of all self-care but unable to carry out any work activities; up and about more than 50% of waking hours
3	Symptomatic and in a chair or in bed for greater than 50% of the day but not bedridden
4	Completely disabled; cannot carry out any self-care; totally confined to bed or chair

Reference: Definition in COSD Core (Performance Status (Adult)), [COSD v10.0 downloads - NDRS](#).